

Triple Inhibitor Mechanism of Antiretroviral from *Sambucus nigra* Phytochemical through Screening Docking

Received 02 June 2023,
Accepted 28 June 2023

DOI: 10.24036/sainstek/vol2-
iss01/18

Nur Sofiatul Aini^{1,2*}, Ahmad Affan Ali Murtadlo^{2,3}, Muhammad Badrut Tamam^{2,4}, Dora Dayu Rahma Turista^{2,5}, Sin War Naw⁶, Md. Emdad Ullah⁷

¹Faculty of Mathematics and Natural Sciences, Universitas Negeri Surabaya, Surabaya, Indonesia

²Generasi Biologi Indonesia Foundation, Gresik, Indonesia

³Faculty of Science and Technology, Universitas Airlangga, Surabaya, Indonesia

⁴Department of Biology, Faculty of Sciences and Technology, Universitas Muhammadiyah Lamongan, Lamongan, Indonesia

⁵Biology Education Department, Faculty of Teacher Training and Education, Mulawarman University, Samarinda, Indonesia

⁶Department of Chemistry, Myitkyina University, Myitkyina, Myanmar

⁷Department of Chemistry, Mississippi State University, Mississippi State, United States

* Corresponding Author: nursofiatul.aini@gmail.com

ABSTRACT

Human immunodeficiency virus type 1 (HIV-1) has a higher infectious nature than HIV-2 because of the genetic drift mechanism. Several HIV-1 strains such as A, B, C, D, E, F, G, H, & J were obtained from Africa, Asia, Europe, Australia, and America. Endemic cases of HIV-1 are often found and HIV-2 is very rare, therefore the development of antiretroviral drugs is currently focused on treating HIV-1 infection⁵. Research on HIV-1 vaccines is still in the research phase and some patients are only taking antiretrovirals to survive. *Sambucus nigra* L. is used by the community for traditional medicine such as influenza A, B, C and SARS-CoV-2 infectious diseases. There are no research reports related to the potential of *Sambucus nigra* L. as an antiretroviral in the treatment of HIV-1 infection and the specific molecular mechanisms that explain it, therefore this research is important for the discovery of alternative antiretroviral drugs from natural materials. The *in silico* method used in this study consists of *Sambucus nigra* L. compounds retrieval, HIV-1 protein preparation, molecular docking, and 3D visualization. *Sambucus nigra* L. can have inhibitory activity on three enzymes of HIV-1 such as protease (PR), reverse transcriptase (RT), integrase (INT) through naringenin and Cyanidin-3,5-diglucoside compounds. Both compounds have more negative binding energy, form stable binding interactions in the target domain, and trigger inhibitory activity.

Keywords: Antiretroviral, HIV-1, *In Silico*, Molecular Docking, *Sambucus nigra*

INTRODUCTION

Human immunodeficiency virus type 1 (HIV-1) from the retrovirus family can trigger opportunistic diseases of the immune system or AIDS^{1,2}. In these conditions the immune system fails due to infection with bacteria,

protozoa, fungi, and other viruses. The identified types of HIV consist of HIV-1 & HIV-2, HIV-1 has a higher infectious nature than HIV-2 because of the genetic drift mechanism^{3,4}. Several HIV-1 strains such as A, B, C, D, E, F, G, H, & J were obtained from Africa, Asia, Europe, Australia, and America^{5,6}. HIV-1 has structural genes consisting of env, gag, and pol then essential elements & regulators such as rev, tat, nef, vpr, vpu, and vif that play a role in helping the virus to evade the immune system's response and pathogenesis in host cells^{7,8}.

† Footnotes relating to the title and/or authors should appear here.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.24036/sainstek/vol2-iss01/18

Endemic cases of HIV-1 are often found and HIV-2 is very rare, therefore the development of antiretroviral drugs is currently focused on treating HIV-1 infection^{9,10}. Research on HIV-1 vaccines is still in the research phase and some patients are only taking antiretrovirals to survive.

HIV-1 has several enzymes consisting of reverse transcriptase (RT), integrase (INT), & protease (PR), all of which play an important role in the replication process^{11,12}. Reverse transcriptase plays a role in the formation of HIV-1 cDNA in several stages consisting of lysyl tRNA having a binding site, RT then adds a nucleotide to synthesize cDNA to the non-coding or U5 and R (repeat) region of the viral RNA^{13,14}. The RT region known as RNase H degrades U5 and R at the 5' end, then tRNA primers synthesize DNA strands. HIV-1 cDNA is carried by integrase to integrate with the host cell genome. Integrase enzymes play a role in the integration of HIV-1 cDNA into the host cell genome and protease enzymes cut viral polypeptides into peptides for the assembly process^{15,16}. RT, INT, and PR activity has a crucial role in viral replication and is very likely to be a target in drug design.

Sambucus nigra L. has health benefits consisting of antidiabetic, anticancer, and antiviral^{17,18}. The content of anthoxyanins in the flowers of *Sambucus nigra* L. is used by the community for traditional medicine such as influenza A, B, and C infectious diseases, the compound works as an antiviral through the mechanism of inhibiting replication activity, immunomodulatory, fever reduction, and anti-inflammatory^{19,20}. *Sambucus nigra* L. was reported to be an alternative treatment in cases of SARS-CoV-2 infection since 2019. There are no research reports related to the potential of *Sambucus nigra* L. as an antiretroviral in the treatment of HIV-1 infection and the specific molecular mechanisms that explain it, therefore this research is important for the discovery of alternative antiretroviral drugs from natural materials.

METHOD

Compound Retrieval

The chemical compounds of *Sambucus nigra* L. ligands in this study consist of Cyanidin-3-sambubioside, Cyanidin-3,5-diglucoside, Catechin, Epicatechin, Naringenin, and Quercetin. PubChem (<https://pubchem.ncbi.nlm.nih.gov/>) is used to retrieve data on ligands such as CID, molecular weight (g/mol), formula, and 3D structure with *structure data format* (sdf) files. Minimization of ligands and conversion to

protein databank (pdb) format were done through OpenBabel software to increase ligand flexibility^{21,22}.

Protein Preparation

This study uses targets from HIV-1 such as protease (PR) for polypeptide cutting, reverse transcriptase (RT) as reverse transcription of cDNA formation, and integrase (INT) which plays a role in the integration of HIV-1 cDNA into the host genome. All target samples were obtained from the RCSB PDB database (<https://www.rcsb.org/>) and then removed water molecules and native ligands through PyMol v2.5 software for optimization and docking preparation^{23,24}.

Molecular Docking

Docking aims to identify the ability of ligand binding to the target by referring to the binding affinity value and interaction pattern on the target domain. The type of docking used in this study is blind to screen the inhibitory activity of *Sambucus nigra* L. on all targets. Autogrid settings are done by covering the entire surface of the target to direct ligand binding which consists of centers and dimensions. Molecular docking in this study was performed by PyRx v.1.0.0 software with Vina Wizard plugin^{25,26}.

3D Visualization

The molecular complex with highest negative binding affinity value was displayed through PyMol v2.5 software with a 3D structure. The visualization method used in this study is structural selection and coloring based on C, H, and O. PyMol plays a role in molecular visualization of three-dimensional structures in proteins and chemical compounds through cartoons, surfaces, sticks, lines, and ribbons^{27,28}.

RESULTS AND DISCUSSION

Information such as name, CID, molecular weight (g/mol), formula, and links on chemical compounds from *Sambucus nigra* L. consisting of Cyanidin-3-sambubioside, Cyanidin-3,5-diglucoside, Catechin, Epicatechin, Naringenin, and Quercetin have been obtained from the PubChem database (Table 1). Previous research revealed that *Sambucus nigra* L. can have potential as an antiviral, antibacterial, antitumor, and antidiabetic, the plant has polyphenol and lectins compounds that are strongly suspected to act as antivirals²⁹. Extracts from *Sambucus nigra* L. can reduce viral load and suppress the spread of viral infection based on in vitro study data.

Table 1. *Sambucus nigra* L. compounds from database

Compound	CID	Molecular Weight (g/mol)	Formula	Link
Cyanidin-3-sambubioside	3084569	616.9	C ₂₆ H ₂₉ CiO ₁₅	https://pubchem.ncbi.nlm.nih.gov/compound/3084569
Cyanidin-3,5-diglucoside	44256718	611.5	C ₂₇ H ₃₁ O ₁₆ ⁺	https://pubchem.ncbi.nlm.nih.gov/compound/44256718
Catechin	9064	290.27	C ₁₅ H ₁₄ O ₆	https://pubchem.ncbi.nlm.nih.gov/compound/9064
Epicatechin	72276	290.27	C ₁₅ H ₁₄ O ₆	https://pubchem.ncbi.nlm.nih.gov/compound/72276
Naringenin	439246	272.25	C ₁₅ H ₁₂ O ₅	https://pubchem.ncbi.nlm.nih.gov/compound/439246
Quercetin	5280343	302.23	C ₁₅ H ₁₀ O ₇	https://pubchem.ncbi.nlm.nih.gov/compound/5280343

Prediction of the binding mechanism of compounds from *Sambucus nigra* on the three HIV-1 proteins was carried out through molecular simulation. The simulation aims to determine the level of binding ability of a ligand to a protein domain based on the binding affinity value in a stable ligand-protein complex and produces negative energy. Binding affinity is formed when there is interaction between protein and ligand, this energy is formed through reversible reactions at constant temperature and pressure in accordance with the laws of thermodynamics^{30,31,32}. The grid in the docking simulation plays a role to direct the ligand binding to the target protein. The simulation performed with refer to a docking grid consisting of Center (Å) X: 15.878 Y: 26.561 Z: 3.595 Dimensions (Å) X: 50.346 Y:

52.213 Z: 59.643 on HIV-1 PR Center (Å) X: 41.297 Y: -0.889 Z: 29.196 Dimensions (Å) X: 112.337 Y: 100.768 Z: 110.694 on HIV-1 RT Center (Å) X: -37.735 Y: 5.417 Z: -2.867 Dimensions (Å) X: 53.962 Y: 45.756 Z: 45.669 on HIV-1 INT. The docking results showed that Naringenin compound has a more negative binding affinity value on PR and Cyanidin-3,5-diglucoside on RT and INT HIV-1 (Table 2). This has answered the molecular mechanism of *Sambucus nigra* medicinal plants can potentially inhibit the activity or infection of the HIV-1 virus. Visualization of the ligand-protein complex structure consisting of Naringenin-PR, Cyanidin-3,5-diglucoside-RT, and Cyanidin-3,5-diglucoside-INT is displayed as sticks and cartoons with staining selection (Figure 1).

Table 2. The results of molecular docking simulation

Compound	CID	Binding Affinity (kcal/mol)		
		PR (PDB ID: 1D4S)	RT (PDB ID: 8DX3)	INT (PDB ID: 6NCJ)
Cyanidin-3-sambubioside	3084569	-7.3	-8.1	-7.0
Cyanidin-3,5-diglucoside	44256718	-7.5	-9.4	-7.8
Catechin	9064	-7.9	-8.2	-6.4
Epicatechin	72276	-8.1	-7.7	-6.5
Naringenin	439246	-8.3	-8.1	-6.5
Quercetin	5280343	-8.2	-8.2	-6.3




Figure 1. Ligand-protein visualization from the docking results. (A) Naringenin-PR (B) Cyanidin-3,5-diglucoside-RT (C) Cyanidin-3,5-diglucoside-INT.

CONCLUSION

Sambucus nigra L. can have inhibitory activity on three enzymes of HIV-1 such as protease (PR), reverse transcriptase (RT), integrase (INT) through naringenin and Cyanidin-3,5-diglucoside compounds. Both compounds have more negative binding energy, form stable binding interactions in the target domain, and trigger inhibitory activity. Further research needs to be conducted such as validation through *in vivo* and *in vitro* approaches for additional scientific evidence.

REFERENCES

- Huerta L. Editorial: Anti-infective 2020: HIV- From pathogenesis to treatment. *Curr Opin Pharmacol.* 2020; 54: x-xii. DOI: 10.1016/j.coph.2020.
- Porter RS, Bode RF. A Review of the Antiviral Properties of Black Elder (*Sambucus nigra* L.) Products. *Phytother Res.* 2017; 31(4): 533-554. DOI: 10.1002/ptr.5782.
- Mahboubi M. *Sambucus nigra* (black elder) as alternative treatment for cold and flu. *ADV TRADIT MED (ADTM).* 2021;21(3):405–14. DOI: 10.1007/s13596-020-00469-z.
- Asgary S, Pouramini A. The Pros and Cons of Using Elderberry (*Sambucus nigra*) for Prevention and Treatment of COVID-19. *Adv Biomed Res.* 2022; 11: 96. DOI: 10.4103/abr.abr_146_21.
- Ramadhani NF, Nugraha AP, Ihsan IS, Agung YA, Rantam FA, Ernawati DS et al. Gingival medicinal signaling cells conditioned medium effect on the osteoclast and osteoblast number in lipopolysaccharide-induced calvaria bone resorption in wistar rats' (*Rattus norvegicus*). *Research Journal of Pharmacy and Technology.* 2021; 14(10): 5232-5237. DOI: 10.52711/0974-360X.2021.00911
- Kharisma VD, Ansori ANM, Jakhmola V, Rizky WC, Widyananda MH, Probojati RT, Murtadlo AAA, Rebezov M, Scherbakov P, Burkov P, Matrosova Y, Romanov A, Sihombing MAEM, Antonius Y, Zainul R. Multi-strain human papillomavirus (HPV) vaccine innovation via computational study: A mini review. *Res J Pharm Technol.* 2022; 15(8):3802-7. doi: 10.52711/0974-360X.2022.00638
- Fahmi M, Kharisma VD, Ansori ANM, Ito M. Retrieval and Investigation of Data on SARS-CoV-2 and COVID-19 Using Bioinformatics Approach. *Adv Exp Med Biol.* 2021; 1318: 839-857. DOI: 10.1007/978-3-030-63761-3_47
- Kharisma VD, Probojati RT, Murtadlo AAA, Ansori ANM, Antonius Y, Tamam MB. Revealing Potency of Bioactive Compounds as Inhibitor of Dengue Virus (DENV) NS2B/NS3 Protease from Sweet Potato (*Ipomoea batatas* L.) Leaves. *Indian J Forensic Med Toxicol.* 2020; 15(1): 1627–1632. DOI: 10.37506/ijfamt.v15i1.13644
- Husen SA, Winarni D, Salamun, Ansori ANM, Susilo RJK, Hayaza S. Hepatoprotective Effect of Gamma-mangostin for Amelioration of Impaired Liver Structure and Function in Streptozotocin-induced Diabetic Mice. *IOP Conference Series: Earth and Environmental Science.* 2019; 217(1): 012031. DOI: 10.1088/1755-1315/217/1/012031
- Turista DDR, Islamy A, Kharisma VD, Ansori ANM. Distribution of COVID-19 and Phylogenetic Tree Construction of SARS-CoV-2 in Indonesia. *J Pure Appl Microbiol.* 2020; 14: 1035-1042. doi: 10.22207/JPAM.14.SPL1.42
- Kharisma VD, Widyananda MH, Ansori ANM, Nege AS, Naw SW, Nugraha AP Tea catechin as antiviral agent via apoptosis agonist and triple inhibitor mechanism against HIV-1 infection: A bioinformatics approach. *J Pharm Pharmacogn Res.* 9(4): 435-445.
- Ansori ANM, Kharishma VD, Muttaqin SS, Antonius Y, Parikesit AA. Genetic Variant of SARS-CoV-2 Isolates in Indonesia: Spike

- Glycoprotein Gene. *J Pure Appl Microbiol.* 2020; 14: 971-978. DOI: 10.22207/JPAM.14.SPL1.35
13. Widyananda MH, Pratama SK, Samoedra RS, Sari FN, Kharisma VD, Ansori ANM, Antonius Y. Molecular docking study of sea urchin (*Arbacia lixula*) peptides as multi-target inhibitor for non-small cell lung cancer (NSCLC) associated proteins. *J Pharm Pharmacogn Res.* 2021; 9(4): 484–496.
 14. Kharisma VD, Ansori ANM. Construction of Epitope-Based Peptide Vaccine Against SARS-CoV-2: Immunoinformatics Study. *J Pure Appl Microbiol.* 2020; 14: 999-1005. DOI: 10.22207/JPAM.14.SPL1.38
 15. Kharisma VD, Ansori ANM, Widyananda MH, Utami SL, Nugraha AP. Molecular simulation: The potency of conserved region on E6 HPV-16 as a binding target of black tea compounds against cervical cancer. *Biochemical and Cellular Archives.* 2020; 20: 2795-2802. DOI: 10.35124/bca.2020.20.S1.2795
 16. Kharisma VD, Agatha A, Ansori ANM, Widyananda MH, Rizky WC, Dings TGA, Derkho M, Lykasova I, Antonius Y, Rosadi I, Zainul R. Herbal combination from *Moringa oleifera* Lam. and *Curcuma longa* L. as SARS-CoV-2 antiviral via dual inhibitor pathway: A viroinformatics approach. *J Pharm Pharmacogn Res.* 2022; 10(1): 138-146. DOI: 10.56499/jppres21.1174_10.1.138
 17. Wijaya RM, Hafidzhah MA, Kharisma VD, Ansori ANM, Parikesit AP. COVID-19 In Silico Drug with *Zingiber officinale* Natural Product Compound Library Targeting the Mpro Protein. *Makara J Sci.* 2021; 25(3): 5. DOI: 10.7454/mss.v25i3.1244
 18. Ansori ANM, Fadholly A, Kharisma VD, Nugraha AP. Therapeutic potential of avian paramyxovirus serotype 1 for cancer therapy. *Biochemical and Cellular Archives.* 2020;20:2827-2832. DOI: 10.35124/bca.2020.20.S1.2827
 19. Prahasanti C, Nugraha AP, Kharisma VD, Ansori ANM, Ridwan RD, Putri TPS et al. Un enfoque bioinformático de la exploración con compuestos de hidroxapatita y polimetilmetacrilato como biomaterial de implantes dentales. *Journal of Pharmacy and Pharmacognosy Research.* 2021; 9(5): 746-754.
 20. Kharisma VD, Ansori ANM, Fadholly A, Sucipto TH. Molecular mechanism of caffeine-aspirin interaction in kopi balur 1 as anti-inflammatory agent: A computational study. *Indian Journal of Forensic Medicine and Toxicology.* 2020; 14(4): 4040-4046. DOI: 10.37506/ijfnt.v14i4.12274
 21. Kharisma VD, Widodo N, Ansori ANM, Nugraha AP. A vaccine candidate of zika virus (ZIKV) from polyvalent conserved b-cell epitope on viral glycoprotein: In silico approach. *Biochemical and Cellular Archives.* 2020;20:2785-2793. DOI: 10.35124/bca.2020.20.S1.2785
 22. Padmi H, Kharisma VD, Ansori ANM, Sibero MT, Widyananda MH, Ullah E, Gumenyuk O, Chylichcova S, Bratishko N, Prasedya ES, Sucipto TH, Zainul R. Macroalgae Bioactive Compounds for the Potential Antiviral of SARS-CoV-2: An In Silico Study. *Journal of Pure and Applied Microbiology.* 2022; 16(2): 1018-1027. DOI: 10.22207/JPAM.16.2.26
 23. Antonius Y, Kharisma VD, Widyananda MH, Ansori ANM, Trinugroho JP, Ullah ME, Naw SW, Jakhmola V, Wahjudi M. Prediction of Aflatoxin-B1 (AFB1) Molecular Mechanism Network and Interaction to Oncoproteins Growth Factor in Hepatocellular Carcinoma. *J Pure Appl Microbiol.* 2022;16(3):1844-1854. doi: 10.22207/JPAM.16.3.29
 24. Dibha AF, Wahyuningsih S, Ansori ANM, Kharisma VD, Widyananda MH, Parikesit AA, Sibero MT, Probojati RT, Murtadlo AAA, Trinugroho JP, Sucipto TH, Turista DDR, Rosadi I, Ullah ME, Jakhmola V, Zainul R. Utilization of Secondary Metabolites in Algae *Kappaphycus alvarezii* as a Breast Cancer Drug with a Computational Method. *Pharmacognosy Journal.* 2022; 14(3): 536-543. DOI: 10.5530/pj.2022.14.68
 25. Młynarczyk K, Walkowiak-Tomczak D, Łysiak GP. Bioactive properties of *Sambucus nigra* L. as a functional ingredient for food and pharmaceutical industry. *J Funct Foods.* 2018; 40: 377-390. DOI: 10.1016/j.jff.2017.11.025.
 26. Aini NS, Ansori ANM, Kharisma VD, Syadzha MF, Widyananda MH, Murtadlo AA, et al. Potential Roles of Purslane (*Portulaca oleracea* L.) as Antimetabolic Syndrome: A Review. *Pharmacognosy Journal.* 2022; 14(3): 710-714. DOI: 10.5530/pj.2022.14.90
 27. Listiyani P, Kharisma VD, Ansori AN, Widyananda MH, Probojati RT, Murtadlo AA, et al. In Silico Phytochemical Compounds Screening of *Allium sativum* Targeting the Mpro

- of SARS-CoV-2. *Pharmacognosy Journal*. 2022; 14(3): 604-609. DOI: 10.5530/pj.2022.14.78
28. Aini NS, Kharisma VD, Widyananda MH, Murtadlo AA, Probojati RT, Turista DD, et al. In Silico Screening of Bioactive Compounds from *Syzygium cumini* L. and *Moringa oleifera* L. Against SARS-CoV-2 via Tetra Inhibitors. *Pharmacognosy Journal*. 2022;14(4):267-272. DOI: 10.5530/pj.2022.14.95
29. Aini NS, Kharisma VD, Widyananda MH, Murtadlo AA, Probojati RT, Turista DD, et al. Bioactive Compounds from Purslane (*Portulaca oleracea* L.) and Star Anise (*Illicium verum* Hook) as SARS-CoV-2 Antiviral Agent via Dual Inhibitor Mechanism: In Silico Approach. *Pharmacognosy Journal*. 2022;14(4):352-357. DOI: 10.5530/pj.2022.14.106
30. Ansori AN, Kharisma VD, Parikesit AA, Dian FA, Probojati RT, Rebezov M, Scherbakov P, Burkov P, Zhdanova G, Mikhalev A, Antonius Y, Pratama MRF, Sumantri NI, Sucipto TH, Zainul R. Bioactive Compounds from Mangosteen (*Garcinia mangostana* L.) as an Antiviral Agent via Dual Inhibitor Mechanism against SARS-CoV-2: An In Silico Approach. *Phcog J*. 2022; 14(1): 85-90. DOI: 10.5530/pj.2022.14.12
31. Wicaksono A, Kharisma VD, Parikesit AA. New Perspectives on Reverse Translation: Brief History and Updates. *Universitas Scientiarum*. 2023; 28(1): 1-20.
32. Ningrum SG, R Sasmita, Kharisma VD. Edible Bird's Nest as Potential Food with Anti-Viral and Anti-Inflammatory Properties Against Covid-19: an in Silico Study. 2023; 11(1): 43-50